Hva er tibialisstimulering?

- Afferent nerve stimulation for pelvic floor dysfunction ML Stoller - International Urogynecology Journal, 1999 - SPRINGER-VERLAG
- Cochrane review https://quintet.no/wp-content/uploads/2021/09/Stewart_et_al-2016-Cochrane_Database_of_Systematic_Reviews.pdf
- Finazzi Agrò E, Campagna A, Sciobica F, et al: Posterior tibial nerve stimulation: is the once-a-week protocol the best option?. Minerva Urol Nefrol. 2005, 57 (2): 119-123.

Virkningsmekanismer

- Tai C, Chen M, Shen B, et al: Irritation induced bladder overactivity is suppressed by tibial nerve stimulation in cats. J Urol. 2011, 186 (1): 326-330. 10.1016/j.juro.2011.04.023.
- Jiang CH, Lindstrom S: Prolonged enhancement of the micturition reflex in the cat by repetitive stimulation of bladder afferents. J Physiol. 1999, 517 (Pt 2): 599-605.
- Danisman A, Kutlu O, Akkaya E, et al: Tibial nerve stimulation diminishes mast cell infiltration in the bladder wall induced by interstitial cystitis urine. Scand J Urol Nephrol. 2007, 41 (2): 98-102. 10.1080/00365590600911233.
- Chang CJ, Huang ST, Hsu K, et al: Electroacupuncture decreases c-fos expression in the spinal cord induced by noxious stimulation of the rat bladder. J Urol. 1998, 160 (6 Pt 1): 2274-2279.
- Finazzi Agrò E, Rocchi C, Pachatz C, et al: Percutaneous tibial nerve stimulation produces effects on brain activity: study on the modifications of the long latency somatosensory evoked potentials. Neurourol Urodyn. 2009, 28 (4): 320-324. 10.1002/nau.20651.

Overaktiv blære referanser

- van Balken MR, Vandoninck V, Gisolf KW, et al: Posterior tibial nerve stimulation as neuromodulative treatment of lower urinary tract dysfunction. J Urol. 2001, 166 (3): 914-918. 10.1016/S0022-5347(05)65863-6.
- Vandoninck V, Van Balken MR, Finazzi Agrò E, et al: Posterior tibial nerve stimulation in the treatment of urge incontinence. Neurourol Urodyn. 2003, 22 (1): 17-23. 10.1002/nau.10036.
- Vandoninck V, van Balken MR, Finazzi Agrò E, et al: Percutaneous tibial nerve stimulation in the treatment of overactive bladder: urodynamic data. Neurourol Urodyn. 2003, 22 (3): 227-232. 10.1002/nau.10111.
- Vandoninck V, van Balken MR, Finazzi Agrò E, et al: Posterior tibial nerve stimulation in the treatment of idiopathic nonobstructive voiding dysfunction. Urology. 2003, 61 (3): 567-572. 10.1016/S0090-4295(02)02378-6.
- van der Pal F, van Balken MR, Heesakkers JP, et al: Correlation between quality of life and voiding variables in patients treated with percutaneous tibial nerve stimulation. BJU Int. 2006, 97 (1): 113-116. 10.1111/j.1464-410X.2006.05860.x.

Urinretensjon:

41 – 100 % vellykket behandling:

* Klingler HC, Pycha A, Schmidbauer J, et al: Use of peripheral neuromodulation of the S3 region for treatment of detrusor overactivity: a urodynamic-based study. Urology. 2000, 56 (5): 766-771. 10.1016/S0090-4295(00)00727-5

Vandoninck V, van Balken MR, Finazzi Agrò E, et al: Posterior tibial nerve stimulation in the treatment of idiopathic nonobstructive voiding dysfunction. Urology. 2003, 61 (3): 567-572. 10.1016/S0090-4295(02)02378-6.

Vandoninck V, Van Balken MR, Finazzi Agrò E, et al: Posterior tibial nerve stimulation in the treatment of voiding dysfunction: Urodynamic data. Neurourol Urodyn. 2004, 23 (3): 246-251. 10.1002/nau.10158.

Nevrogene blæreforstyrrelser:

- Amarenco G, Ismael SS, Even-Schneider A, et al: Urodynamic effect of acute transcutaneous posterior tibial nerve stimulation in overactive bladder. J Urol. 2003, 169 (6): 2210-2215. 10.1097/01.ju.0000067446.17576.bd.
- Kabay SC, Kabay S, Yucel M, et al: Acute urodynamic effects of percutaneous posterior tibial nerve stimulation on neurogenic detrusor overactivity in patients with Parkinson's disease. Neurourol Urodyn. 2009, 28 (1): 62-67. 10.1002/nau.20593.
- Kabay S, Kabay SC, Yucel M, et al: The clinical and urodynamic results of a 3-month percutaneous posterior tibial nerve stimulation treatment in patients with multiple sclerosis-related neurogenic bladder dysfunction. Neurourol Urodyn. 2009, 28 (8): 964-968. 10.1002/nau.20733.
- Gobbi C, Digesu G, Khullar V, El Neil S, Caccia G, Zecca C: Percutaneous posterior tibial nerve stimulation as an effective treatment of refractory lower urinary tract symptoms in patients with multiple sclerosis: preliminary data from a multicentre, prospective, open label trial. Mult Scler. 2011, 17 (12): 1514-1519. 10.1177/1352458511414040. Epub 2011 Jul 14

Effekt CPP:

- van Balken MR, Vandoninck V, Messelink BJ, et al: Percutaneous tibial nerve stimulation as neuromodulative treatment of chronic pelvic pain. Eur Urol. 2003, 43 (2): 158-163. 10.1016/S0302-2838(02)00552-3.
- Kim SW, Paick JS, Ku JH: Percutaneous posterior tibial nerve stimulation in patients with chronic pelvic pain: a preliminary study. Urol Int. 2007, 78 (1): 58-62. 10.1159/000096936.
- Kabay S, Kabay SC, Yucel M, et al: Efficiency of posterior tibial nerve stimulation in category IIIB chronic prostatitis/chronic pelvic pain: a Sham-Controlled Comparative Study. Urol Int. 2009, 83 (1): 33-38. 10.1159/000224865.
- Zhao J, Nordling J: Posterior tibial nerve stimulation in patients with intractable interstitial cystitis. BJU Int. 2004, 94 (1): 101-104. 10.1111/j.1464-410X.2004.04909.x.

- Zhao J, Bai J, Zhou Y, et al: Posterior tibial nerve stimulation twice a week in patients with interstitial cystitis. Urology. 2008, 71 (6): 1080-1084. 10.1016/j.urology.2008.01.018.
- Baykal K, Senkul T, Sen B, et al: Intravesical heparin and peripheral neuromodulation on interstitial cystitis. Urol Int. 2005, 74 (4): 361-364. 10.1159/000084439.
- Congregado Ruiz B, Pena Outeiriño XM, Campoy Martínez P, León Dueñas E, Leal López A: Peripheral afferent nerve stimulation for treatment of lower urinary tract irritative symptoms. Eur Urol. 2004, 45: 65-69. 10.1016/j.eururo.2003.08.012.
- Gokyldiz S, et al: Effects of percutaneous tibial nerve stimulation therapy on chronic pelvic pain. Gynecol Obstet Invest. 2012, 73 (2): 99-105. 10.1159/000328447. Epub 2012 Jan 20

Ikke nevrogene blæreforstyrrelser hos barn:

- Hoebeke P, Renson C, Petillon L, et al: Percutaneous electrical nerve stimulation in children with therapy resistant nonneuropathic bladder sphincter dysfunction: a pilot study. J Urol. 2002, 168 (6): 2605-2607. 10.1016/S0022-5347(05)64227-9.
- De Gennaro M, Capitanucci ML, Mastracci P, et al: Percutaneous tibial nerve neuromodulation is well tolerated in children and effective for treating refractory vesical dysfunction. J Urol. 2004, 171 (5): 1911-1913. 10.1097/01.ju.0000119961.58222.86.
- Capitanucci ML, Camanni D, Demelas F, et al: Long-term efficacy of percutaneous tibial nerve stimulation for different types of lower urinary tract dysfunction in children. J Urol. 2009, 182 (Suppl 4): 2056-2061.
- Burton C, Sajja A, Latthe PM. Efectiveness of percutaneous posterior tibial nerve stimulation for overactive bladder: a systematic review and meta-analysis. Neurourol Urodyn. 2012. https://doi.org/10.1002/nau.22251.
- **Neuromodulation for Treatment of Refractory Non-neurogenic Lower Urinary Tract Symptoms in Children: an Overview**Tong, Ching Man Carmen ; Kitchens, David M.Current bladder dysfunction reports, 2023-03, Vol.18 (1), p.59-6

Obstipasjon:

- Gokce AH, Gokce FS, Iliaz R, Gulaydin N. Transcutaneous Tibial Nerve Stimulation as Therapy for Functional Constipation. Turk J Gastroenterol. 2022 Jul;33(7):565-569. doi: 10.5152/tjg.2022.21342. PMID: 35879913; PMCID: PMC9404683.
- Velasco-Benitez C, Villamarin E, Mendez M, Linero A, Hungria G, Saps M. Efficacy of transcutaneous posterior tibial nerve stimulation in functional constipation. Eur J Pediatr. 2023 Mar;182(3):1309-1315. doi: 10.1007/s00431-022-04798-w. Epub 2023 Jan 13. PMID: 36637541; PMCID: PMC9838342.

Analfissurer:

- https://quintet.no/forskning/tibialisstimulering-ved-analfissurer/
-

Obstipasjon:

- Transcutaneous Electrical Nerve Stimulation to Improve Female Sexual Dysfunction Symptoms: A Pilot Study **Lauren L. Zimmerman MS, Priyanka Gupta MD, Florence O'Gara BSN, MSW, Nicholas B. Langhals PhD, Mitchell B. Berger MD, PhD, Tim M. Bruns PhD**